

Questionnaire for CT Appointment CT検査予約問診票

<small>フリガナ</small>		Gender 性別	Age 年齢	Height 身長	Weight 体重
Patient Name <small>氏名</small>		Male / Female <small>男性 女性</small>		cm	kg
Purpose of Examination <small>検査目的</small>					

Type of exam <small>検査内容</small>	<input type="checkbox"/> Simple Study <small>単純検査</small> <input type="checkbox"/> Contrast study (See "Particular Comments" and check applicable <input type="checkbox"/>) <small>造影検査</small>				
Imaging parts <small>撮影部位</small>	<input type="checkbox"/> Brain <small>脳</small>	<input type="checkbox"/> Cervix <small>頸部</small>	<input type="checkbox"/> Chest <small>胸部</small>	<input type="checkbox"/> Abdomen(liver - kidney) <small>腹部</small>	<input type="checkbox"/> Cavitas pelvis <small>骨盤腔</small>
	<input type="checkbox"/> Orbit <small>眼窩</small>	<input type="checkbox"/> Pranasal sinuses <small>副鼻腔</small>	<input type="checkbox"/> Cervical vertebrae <small>頸椎</small>	<input type="checkbox"/> Thoracic vertebrae <small>胸椎</small>	<input type="checkbox"/> Lumbar vertebrae <small>腰椎</small>
	<input type="checkbox"/> Others <small>その他</small> ()				

Output images <small>画像出力</small>	<input type="checkbox"/> No need <small>不要</small> <input type="checkbox"/> Need <small>要</small> → If need, how? <small>要の場合選択</small> : <input type="checkbox"/> By CD <small>CD出力</small> <input type="checkbox"/> By films <small>フィルム出力</small>			
---------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--	--	--

Pick up result <small>結果渡し *No mailing service</small>	<input type="checkbox"/> The patient picks up on the same day (takes 1-3 hours) <small>当日日本人手渡し</small> <input type="checkbox"/> The patient picks up another day. <small>後日、本人来院</small> <input type="checkbox"/> A staff from referring facility picks up another day. <small>後日紹介元スタッフ来院</small>			
------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--	--	--

Implanted medical device <small>体内チェック事項</small>	Pacemaker <small>心臓ペースメーカー</small> :	<input type="checkbox"/> Yes	<input type="checkbox"/> No
	Implantable cardioverter defibrillator(ICD) <small>槽込み型除細動器</small> :	<input type="checkbox"/> Yes	<input type="checkbox"/> No

Particular comments for contrast exam <small>造影特記事項</small> *Please study the notes below and check all applicable <input type="checkbox"/> .	1) Have you ever had any contrast exams in the past? <small>造影の既往</small> <input type="checkbox"/> Yes <input type="checkbox"/> No 2) Are you allergic to anything? <small>アレルギーの既往</small> <input type="checkbox"/> Yes <input type="checkbox"/> No 3) Are you allergic to contrast medium? <small>造影剤アレルギー</small> <input type="checkbox"/> Yes <input type="checkbox"/> No 4) Have you ever been diagnosed with asthma? <small>ぜんそくの既往</small> <input type="checkbox"/> Yes <input type="checkbox"/> No <small>*If yes, no contrast but simple study only.</small> 5) Have you ever been diagnosed with heart disease? <small>心疾患既往</small> <input type="checkbox"/> Yes <input type="checkbox"/> No 6) Are you currently taking any hypoglycemic drugs? <small>血糖降下剤の内服</small> <input type="checkbox"/> Yes <input type="checkbox"/> No ※ If yes, the patient is explained enough to understand that... <small>有の場合、</small> <input type="checkbox"/> : it is necessary to stop taking the drugs 48 hours before and after using some contrast medium, if his/ her kidney function is decreasing. <small>腎機能が低下している場合は造影剤使用前48時間、使用后48時間休薬の必要があることの説明</small> <input type="checkbox"/> : if the patient did not stop taking drugs 48 hours before, it would be only a simple study, no contrast study performed. <small>休薬がなされていない場合造影検査は施行できず、単純CTのみ施行可となることの説明</small> 7) Do you currently have a central venous port implanted which is safe for the contrast study? <small>造影可能な埋込み型中心静脈CVポート</small> <input type="checkbox"/> Yes <input type="checkbox"/> No 8) <input type="checkbox"/> : Data of Creatinine within 3 months.(within a few days if possible) <small>3ヶ月以内のクレアチニンの検査データ(できれば数日以内)</small> Level値 : (Date of test検査日:) <input type="checkbox"/> Currently testing <small>現在検査中</small> 9) <input type="checkbox"/> : eGFR(estimated glomerular filtration rate) is over 50. <small>eGFR(推定糸球体濾過値)が50を超えていることの確認</small> <small>*If rate was under 50, it would be a simple study, no contrast study performed. 50以下の場合、造影検査は施行できず、単純CTのみ施行可</small> 10) <input type="checkbox"/> : The patient is explained enough to understand to have some contrast medium in. <small>患者へ造影剤を使用することを説明済</small>		
------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--	--

【NOTES】 造影検査の注意点

3) 4) → At our Radiology department, we do not perform any contrast studies to the patient who is allergic to contrast medium or have asthma, but simple MRI only.

6) → There is a critical risk on the side effect when some kinds of hypoglycemic drug (Biguanide diabetes drug) is taken along with some iodinated contrast medium. Patients are required to stop taking drug 48 hours before and after having contrast medium in, if their eGFR levels are under 60 and they are taking drug shown as below. If the patient did not stop taking drugs in advance, we do not perform any contrast studies but simple CT only.

Metglucoメトグルコ錠 ・ Metformin Hydrochlorideメトホルミン塩酸塩錠 ・ Glyceranグリコラン錠 ・ Nelbisネルビス錠 ・ Medetメデット錠 ・ Metact Combination Tabletsメタクト配合錠
 EquMet Combination Tablets エクメット配合錠 ・ INISYNC Combination Tabletsイニシンク配合錠 ・ Dibetosジベトス錠 ・ Dibeton S Enteric coated Tabletsジベトンス腸溶錠

8) 9) → The Creatinine data within 3 months (within a few days if possible) is required as we must know that the kidney function is not decreasing before the contrast CT. If eGFR rate was under 50, we perform a simple study without contrast. For the patients who has been on dialysis, we perform contrast CT as usual.